Contents lists available at ScienceDirect





journal homepage: www.elsevier.com/locate/yqres

Quaternary Research

Changes in the glacier extent and surface elevation along the Ningchan and Shuiguan river source, eastern Qilian Mountains, China



Bo Cao ^a, Baotian Pan ^{a,*}, Jie Wang ^a, Donghui Shangguan ^b, Zhenling Wen ^a, Wentao Qi ^a, Hang Cui ^a, Yaoyang Lu ^a

^a MOE Key Laboratory of Western China's Environmental Systems, Collaborative Innovation Centre for Arid Environments and Climate Change, Lanzhou University, Gansu, Lanzhou 730000, China ^b State Key Laboratory of Cryospheric Science, Cold and Arid Regions Environmental and Engineering Research Institute, Chinese Academy of Sciences, Lanzhou 730000, China

ARTICLE INFO

Article history: Received 6 August 2013 Available online 25 February 2014

Keywords: GPS RTK Glacier extent Surface elevation Qilian Mountains

ABSTRACT

We investigate the changes at nine glaciers in the Ningchan and Shuiguan river source, eastern Qilian Mountains, between 1972 and 2010. According to analysis of topographic maps and multispectral satellite data, all nine glaciers in the study area have retreated, by a maximum of 250 ± 57.4 m and a minimum of 91 ± 57.4 m. The total glacier area decreased by 1.20 km^2 , corresponding to 9.9% of the glacierized area in 1972. Comparing the two DEMs generated from the topographic maps and Real-Time Kinematic GPS data, the mean glacier thinning rate was 0.64 m yr^{-1} between 1972 and 2010. The most significant thinning generally occurred on the termini. The ice-volume loss was about $106.8 \pm 46.7 \times 10^{-3} \text{ km}^3$ (equal to $90.8 \pm 39.7 \times 10^{-3} \text{ km}^3$ w.e.), which suggested a mean water discharge of $0.1 \pm 0.05 \text{ m}^3$ /s during 1972–2010. Based on analysis of meteorological data, the summer temperature (June–August) tends to increase over a similar time period. The consistency of temperature increase and glacier shrinkage allows us to suggest that air temperature plays an important role in glacier changes in this region.

© 2014 University of Washington. Published by Elsevier Inc. All rights reserved.

Introduction

Melting of glaciers and polar ice caps is believed to significantly raise sea levels (Kaser et al., 2006; Meier et al., 2007; Gardner et al., 2013). To understand the changes in glacier length, areal extent and mass balance is thus important for reconstructing past climates and evaluating the contribution of glaciers to modern sea levels (Oerlemans, 2005; Meier et al., 2007; Ohmura, 2011). The key to understanding the mechanisms of glacier variation likely lies in a thorough analysis of glacier response to climate change (Lemke, 2007; Anderson et al., 2008). Glacier meltwater also plays an important role as a source of freshwater in arid and semi-arid areas, where it supports daily life and the development of agriculture, livestock and industry (Yao et al., 2004; Ding et al., 2006; Kaser et al., 2010). Mass loss of glaciers has wide-ranging consequences such as changing runoff distribution (Gao et al., 2012; Zhang et al., 2012), sea-level rise (Radić and Hock, 2011; Jacob et al., 2012) and increasing risk of glacial lake outburst floods (Bolch et al., 2011; Xie et al., 2013).

It has been suggested that the mid-latitude Asian glaciers have retreated continuously since the 1960s as a result of global warming (Liu et al., 2003, 2006; Shangguan et al., 2006; Aizen et al., 2007; Bolch, 2007; Gardner et al., 2013). In addition, Yao et al. (2012) reported ten of the 15 glaciers in the Tibetan Plateau (TP) exhibited a negative mass balance, the most negative being about 1.7 m yr⁻¹. Based on the

* Corresponding author. *E-mail address:* panbt@lzu.edu.cn (B. Pan). mass balance measurements of the Qivi, Small Dongkemadi and Kangwure glaciers, the glaciers have shown an accelerating trend of more negative mass balance in recent years (Yao et al., 2012), as also have the Akshiirak glaciers, Tien Shan, Central Asia (Surazakov and Aizen, 2006). However, due to the remote location and wide distribution of these glaciers, ice thickness and ice volume monitoring is sparse. Traditional ground-based techniques have played an essential role in measuring glacier mass balance directly, but they tend to be laborious, expensive and provide very limited spatial coverage (Keutterling and Thomas, 2006; Racoviteanu et al., 2008). Fortunately, studies of glaciers have improved due to novel, rapid and relatively inexpensive techniques such as remote sensing and aerial photography (Paul et al., 2004; Liu et al., 2006), GIS and GPS technologies (Kargel et al., 2005). Several studies have used digital elevation models (DEMs) based on remotely sensed data and GPS technologies to measure changes in glacier surface elevation (e.g. Rivera and Casassa, 1999; Rignot et al., 2003; Berthier et al., 2004; King, 2004; Rivera et al., 2005; Howat et al., 2007; Berthier and Toutin, 2008).

Monitoring the change in glacier area using remote-sensing data indicates that glaciers in the Qilian Mountains have tended to retreat during the past 50 yr (Liu et al., 2003; Ding et al., 2006; Shangguan et al., 2010). Glaciers in the Lenglongling Mountains, eastern Qilian Mountains, are small and their melt will not contribute significantly to global sea level rise. However, they are sensitive indicators of climatic change at a relatively short time scale (a few decades), and they serve as benchmark glaciers for many global applications. Their disappearance might have great economic and societal impacts, for example on

0033-5894/\$ – see front matter © 2014 University of Washington. Published by Elsevier Inc. All rights reserved. http://dx.doi.org/10.1016/j.yqres.2014.01.011 the hydrologic regime and tourism in the Hexi Corridor. Pan et al. (2012) determined glacier area change in the western Lenglongling Mountains. Xie et al. (1985) and Liu et al. (1992) showed some glacier termini retreated in the Lenglongling Mountains from 1956 to 1984. Li et al. (2010) estimated an average thinning of 15 ± 8 m for Shuiguan (SG) river source glacier No. 4 using field measurements in 2007 and topographical map data from 1972. These studies are very important for understanding the response of glacier mass balance to climatic change in these areas. However, more data are needed to make a robust estimation. To update research on glacier mass balance in this region, we chose nine glaciers in the Ningchan (NC) and SG river source, which originate in the Lenglongling Mountains. The aim of this study is to gain a better understanding of glacier mass balance in terms of length, area and ice-surface elevation change from 1972 to 2010.

Study location

The Lenglongling Mountains are located in the northeastern part of the TP (Fig. 1a). The Chinese Glacier Inventory (CGI) (Wang et al., 1981) shows that a total of 244 glaciers occur above 4000 m asl in the Lenglongling Mountains, covering a total area of 103.02 km² (Fig. 1b). Glaciers in this area are mainly influenced by the combined effects of the East Asian monsoon and the Westerlies (Chen et al., 2008), and are classified as subcontinental-type glaciers (Shi and Liu, 2000). Mean annual air temperature in the period 1980–2010 was about 1.17°C at Menyuan meteorological station (37°23'N, 101°37'E, 2924 m asl). The mean equilibrium line attitude (ELA) in the Lenglongling Mountains is about 4500 m (Wang et al., 1981) and, extrapolating with a standard atmospheric lapse rate of 0.72°C (100 m)⁻¹ (Wang

et al., 2009), mean annual air temperature at the ELA is about -10.18 °C (Pan et al., 2012).

There are ten glaciers in the NC and SG river source, located on the northern slope of the Lenglongling Mountains. The highest peak has an elevation of 5024 m asl. One of the glaciers is a hanging glacier with a terminus altitude of 4580 m asl. All the other glacier terminus altitudes are about 4100–4200 m asl (Wang et al., 1981); the area of these nine glaciers varies and the average area is 1.34 km². The largest glacier covers more than 3 km², and the smallest less than 0.5 km² (Fig. 1c).

Data and methods

GPS data

GPS measurement has been applied in glacier research for many years (King, 2004; Elizabeth et al., 2011; Kunz et al., 2012). A GPS-RTK (Real-Time Kinematic) survey of glaciers in NC and SG river source was conducted from 12 to 20 July, 2010 (Table 1) (Fig. 2a). Five Trimble 5800 dual-frequency GPS receivers were used to implement the RTK survey. The points shown in Figure 2a were obtained in RTK mode, with four receivers carried by four researchers, and the fifth receiver (used as the base station for differential correction of post-processing) installed on an outcrop. The survey areas shown in Figure 2 are accessible for fieldwork. The GPS-measured area covers about 38.8% of the total area in 2010.

The precisions of the vertical and horizontal measurements by GPS were about 0.1 m and 0.05 m, respectively. In glacierized areas, the precision may decrease for unavoidable reasons, such as clouds or



Figure 1. Location of the Ningchan and Shuiguan river source glaciers. (a) The inset shows the location of the study area within the TP. Dashed line shows the boundary between westerly and monsoon moisture (after Chen et al., 2008). (b) Overview of the study area overlaid on DEM. (c) Overview of the borderlines of the glacier with a Landsat image. NC01 stands for Ningchan river source glacier No. 01 and SG01 stands for Shuiguan river source glacier No. 01.

Table 1

source of data used in this study.							
Data source	Date	Resolution	RMSE ^b	DEM error			
Topographic map	1972	5 m/1:50,000 ^a		<11 m			
Landsat TM	11 Jun 1995	28.5 m	<28.5 m				
Landsat TM	08 Sep 2010	28.5 m	<28.5 m				
GPS measurements	2010	0.3 m		0.5 m			

^a According to the criteria of the State Bureau of Surveying and Mapping of China, the ground resolution of maps at scales of 1:50,000 is about 5 m.

^b The RMSE of remote-sensing images is <1 pixel.

ionosphere error. However, the vertical accuracy is less than 0.3 m and horizontal accuracy is less than 0.1 m (Rivera et al., 2005). All GPS point data were re-projected to the Universal Transverse Mercator (UTM) coordinate system and referenced to the World Geodetic System 1984 (WGS84) and then the point data were interpolated by the Inverse Distance Weighted (IDW) method to produce the raster of DEM2010, with a grid resolution of 15 m (Fig. 2b).

Aerial photograph and topographic maps

We used a map derived from aerial photographs at a scale of 1:50,000 acquired in 1972 by the State Bureau of Surveying and Mapping. Then we digitized the 20 m interval contours and spot heights in order to generate a Triangular Irregular Networks (TIN) map and create a DEM1972 based on Beijing54. The center and flattening of Beijing54 are different from the WGS84 UTM. In order to compare with other data, the DEM was re-projected to the WGS84 UTM datum using a seven-parameter datum transformation model with a 15 m resolution. The application of the seven-parameter datum transformation model has been discussed by Wang et al. (2003) and Zhang et al. (2010). The error using a seven-parameter datum transformation model is <0.002



Figure 2. GPS-RTK survey in the study region. (a) Map of the surface points. (b) DEM generated by GPS survey data.

m (Wang et al., 2003). The identified glacier outline was then digitized using GIS based on vector files.

Remote-sensing data

Two Landsat Thematic Mapper (TM) images (Path: 132, Row: 34, 11 June 1995; 08 September 2010) were obtained to identify the glacier borderline. Although some areas in the TM image 2010 were covered with seasonal snow, NC and SG river source glaciers can be distinguished clearly from other surface types (Pan et al., 2012). The Landsat TM data are obtained from the Global Land Cover Facility (GLCF) and the United States Geological Survey (USGS). They were all orthorectified to the WGS84 UTM datum. The best tool to extract more reliable information from satellite images for many glaciers is manual digitization (Raup et al., 2007).

Surface elevation changes

The lower parts of each glacier were selected to calculate glacier surface elevation changes between 1972 and 2010 (Fig. 2a). Surface elevation or ice volume changes were derived from repeated comparisons of the DEMs for different years. Ice volume changes were calculated by multiplying the mean value of surface elevation change, ΔH , by the glacier surface area, *S*.

Results

Changes in glacier terminus position and area

All nine glaciers retreated during the period 1972–2010, with a mean rate of 4.7 m yr⁻¹. The maximum and minimum values were 250 ± 57.4 m and 91 ± 57.4 m, respectively (Table 2). The maximum glacier length retreat occurred on NC04 glacier. In addition, several field campaigns to study the glaciers in this region have been carried out since 1956. These studies show glacier retreat of about 400 m (20 m yr⁻¹) during 1956–76 (Xie et al., 1985) and 71.2 m (8.9 m yr⁻¹) during 1976–84 (Liu et al., 1992). The glaciers have shrunk continuously since at least 1956.

Glacier terminus altitude rises as the glacier shrinks. The rise is rapid when the glacier terminus is steep, and slow when it is flat. The glacier terminus altitudes rose from 2 m (NC06) to 81 m (NC04) from 1972 to 2010 for nine glaciers (Table 2). The increase in mean terminus altitude is 39.3 m in NC and SG river source glaciers. The ELA has increased more than 100 m since 1972 as evidenced by field observation in 2007 (Li et al., 2010).

The total area decrease for these nine glaciers was 1.20 km², accounting for 9.9% of the total area in 1972 (Table 3). As each glacier has a different shape, the glacier retreat rates varied significantly. Some glaciers shrunk relatively faster, e.g. NC01 ($8.2 \times 10^{-3} \text{ km}^2 \text{ yr}^{-1}$) and SG03 ($6.1 \times 10^{-3} \text{ km}^2 \text{ yr}^{-1}$), while others retreated slowly, e.g. NC04 ($0.8 \times 10^{-3} \text{ km}^2 \text{ yr}^{-1}$) and SG04 ($1.8 \times 10^{-3} \text{ km}^2 \text{ yr}^{-1}$).

Changes in glacier surface elevation and ice volume

The glaciers thinned by between 2 and 70 m between 1972 and 2010 (Fig. 3a). The trend of surface elevation change fluctuated with altitude. The mean elevation change was 24.5 m (0.64 m yr⁻¹) for nine glaciers for all surveyed areas (Table 4). The largest ice thinning was observed at NC01 glacier, 0.76 m yr⁻¹ from 1972 to 2010. Generally, most thinning occurred in the lower sections near the terminus area for 35 m in the study area (below 4250 m asl) (Fig. 3b). The thinning trend appears to decrease at higher altitudes.

Table 4 shows the ice volume loss of each glacier from 1972 to 2010. The total ice volume loss was 106.8×10^{-3} km³, corresponding to 90.8×10^{-3} km³ w.e. if the ice density is assumed to be 0.85×10^3 kg m⁻³ (Huss, 2013).

B. Cao et al. / Quaternary Research 81 (2014) 531-537

Table 2

Changes in glacier terminus.

Glaciers		NC				SG				
		01	02	03	04	06	01	02	03	04
Terminus altitude (m)	1972 2010	$\begin{array}{c} 4170 \pm 11 \\ 4225 \pm 0.3 \end{array}$	$\begin{array}{c} 4175\pm11\\ 4237\pm0.3\end{array}$	$\begin{array}{c} 4140\pm11\\ 4156\pm0.3\end{array}$	$\begin{array}{c} 4160\pm11\\ 4241\pm0.3\end{array}$	$\begin{array}{c} 4158\pm11\\ 4160\pm0.3\end{array}$	$\begin{array}{c} 4158\pm11\\ 4205\pm0.3\end{array}$	$\begin{array}{c} 4117\pm11\\ 4130\pm0.3\end{array}$	$\begin{array}{c} 4172\pm11\\ 4198\pm0.3\end{array}$	$\begin{array}{c} 4193\pm11\\ 4245\pm0.3\end{array}$
Terminus change (m)	1956–1976 1976–1984			400 ^a	400 ^a	200 ^a	310 ^a	450 ^a	360 ^a	320 ^a 71.2 ^b
	1972-1995	91	124	56	176	85	152	145	140	201
	1995-2010	79	40	35	74	54	51	32	50	55
	1972-2010	170	164	91	250	139	203	177	190	232
Terminus retreat rate (m yr^{-1})	1956–1976 1976–1984			20.0	20.0		15.5	22.5	18.0	16.0 ^a 8.9 ^b
	1972-1995	4.0	5.4	2.4	7.7	3.7	6.6	6.3	6.1	8.7
	1995-2010	5.3	2.7	2.3	5.0	3.6	3.4	2.1	3.3	3.7
	1972-2010	4.5	4.3	2.4	6.6	3.6	5.3	4.6	5.0	6.1

^a After Xie et al. (1985). ^b After Liu et al. (1992).

However, those parts of the glaciers that were lost due to terminus retreat were excluded from these ice volumes. Furthermore, glacier surface elevation changes were not determined for the higheraltitude areas of the glaciers. So, the ice volume loss from 1972 to 2010 was more than 106.8×10^{-3} km³ in the NC and SG river source glaciers.

Discussion

Analysis of errors

The accuracy in measuring glacier terminus positions extracted from multi-temporal satellite images primarily relies on the sensor resolutions (Williams et al., 1997; Racoviteanu et al., 2009; Paul et al., 2013) and the co-registration errors (Hall et al., 2003; Silverio and Jaquet, 2005). According to Ye et al. (2006) and Silverio and Jaquet (2005), the measurement uncertainty of the glacier terminus, U_T , can be calculated from

$$U_T = \sqrt{\Sigma \lambda^2} + \sqrt{\Sigma \varepsilon^2} \tag{1}$$

where U_T is the measurement uncertainty of the glacier terminus in the study area, λ is the original pixel resolution of each image and ϵ is the co-registration error (<1 pixel) of each image to the topographic map generated in 1972. For the images in 1972, 1995 and 2010, the resolutions are 5 m, 28.5 m and 28.5 m respectively. According to Eq. (1), the uncertainties in measurement of the glacier terminus retreat were 57.4 m during the periods 1972-95 and 1972-2010, and 68.8 m

between 1995 and 2010. The uncertainty in glacier area fron	ı 1972	to
2010 is calculated to be 0.004 km ² from		

$$U_A = 2U_T \sqrt{\Sigma \lambda^2} + \Sigma \varepsilon^2 \tag{2}$$

where U_A is the uncertainty in calculating glacier extent.

The vertical errors were generated from the DEM1972 and DEM2010. DEM1972 is the main source of systematic error in our analysis. The systematic vertical errors of the DEM generated from the 1:50,000 topographic maps were less than 11 m on slopes less than 15° (State Bureau of Surveying and Mapping, 2007). In order to calculate the accuracy error of the DEM1972, some points on the stable terrain were surveyed and the error was about 5 m in the vertical (Table 5). However, the points are too few to have statistical significance, so we use the maximum error of 11 m as the vertical error. The accuracy of the GPS surveys is 0.3 m; however, when we use these points to create a DEM, the error for the 2010DEM will be larger than 0.3 m, which was determined by the number of points and whether the distribution was uniform. Therefore we randomly chose 30 points from each glacier to examine the 2010DEM. The survey points are very dense and the glacier surface is relatively smooth in the study area. The mean elevation error was small, about 0.5 m.

The vertical error of these two DEMs was about 11 m as calculated by Eq. (3) (Barrand et al., 2010)

$$Error = \sqrt{Error^2}_{\text{DEM 1972}} + Error^2_{\text{DEM 2010}} = \sqrt{11^2 + 0.5^2} \approx 11 \text{ m.} \quad (3)$$

Changes in glacier area.

Glaciers		NC					SG				
		01	02	03	04	06	01	02	03	04	
Area (km ²)	1972	0.78	0.36	1.54	1.28	1.22	0.52	3.15	1.47	1.82	
	1995	0.60	0.29	1.47	1.26	1.15	0.46	3.05	1.32	1.78	
	2010	0.47	0.25	1.43	1.25	1.12	0.43	3.00	1.24	1.75	
	GPS measured	0.34	0.11	0.52	0.37	0.13	0.25	1.24	0.77	0.52	
Area change (km ²)	1972-1995	0.18	0.07	0.07	0.02	0.07	0.06	0.10	0.15	0.04	
	1995-2010	0.13	0.04	0.04	0.01	0.03	0.03	0.05	0.08	0.03	
	1972-2010	0.31	0.11	0.11	0.03	0.10	0.09	0.15	0.23	0.07	
Area retreat rate ($\times 10^{-3}$ km ² yr ⁻¹)	1972-1995	7.8	3.0	3.0	0.9	3.0	2.6	4.3	6.5	1.7	
	1995-2010	8.7	2.7	2.7	0.7	2.0	2.0	3.3	5.3	2.0	
	1972-2010	8.2	2.9	2.9	0.8	2.6	2.4	3.9	6.1	1.8	

B. Cao et al. / Quaternary Research 81 (2014) 531-537



Figure 3. Glacier surface elevation change. (a) Changes in glacier surface thickness. (b) Ice thickness and slope changes along the paths on each glacier.

Reasons for glacier variation

Precipitation and air temperature are the two main factors controlling glacier changes (Oerlemans, 2005). Precipitation determined the glacier's accumulation, while summer temperature determined the melt. Menyuan meteorological station and Xining radiosonde station (36°43'N, 101°45'E, 2295 m asl) are about 20 km and 100 km south of the study area, respectively. The meteorological data show that the annual precipitation presents a fluctuation but with no significant trend. However, the summer air temperature and freezing-level height (FLH) show a dramatic increase (Fig. 4) which causes high summer melt rates and negative glacier mass balance. Therefore, the glacier variations in the NC and SG river source appear to be concurrent with changes of summer air temperature and FLH during the period 1972–2010. Pan et al. (2012) pointed out that glaciers in the Western Lenglongling Mountains shrunk between 1972 and 2007, and the glacier shrinkage

Table 4

Glacier surface elevation and volume changes in GPS measured.

				NC				SG			
01	02	03	04	06	01	02	03	04			
Area $(\%)^a$ 72 Mean slope (°) 13.8 Elevation bands (m) 4240-4 Mean surface elevation change (m) 28.8 ± Mean thinning rate (m a ⁻¹) 0.76 ±	$\begin{array}{r} 44\\ 17.6\\ 508 \\ 4237-4350\\ 11 \\ 23.7 \pm 11\\ 0.29 \\ 0.62 \pm 0.29\\ \end{array}$	$\begin{array}{c} 36 \\ 15.1 \\ 4151-4602 \\ 28.4 \pm 11 \\ 0.75 \pm 0.29 \end{array}$	$\begin{array}{c} 30 \\ 15.7 \\ 4166 - 4520 \\ 22.0 \pm 11 \\ 0.58 \pm 0.29 \end{array}$	$12 \\ 21.7 \\ 4157-4337 \\ 22.0 \pm 11 \\ 0.58 \pm 0.29 \\ 0.000 \\ 0.$	$58 \\ 21.6 \\ 4216-4500 \\ 24.0 \pm 11 \\ 0.63 \pm 0.29 \\ 100000000000000000000000000000000000$	$\begin{array}{c} 41 \\ 14.2 \\ 4150-4550 \\ 27.5 \pm 11 \\ 0.72 \pm 0.29 \end{array}$	$\begin{array}{c} 62 \\ 16.9 \\ 4200-4678 \\ 22.7 \pm 11 \\ 0.60 \pm 0.29 \end{array}$	$\begin{array}{c} 30 \\ 17.1 \\ 4250-4644 \\ 21.4 \pm 11 \\ 0.56 \pm 0.29 \end{array}$			

^a GPS measured area covers the percentage of the total area in 2010.

_	2	\sim
5		h
-	-	υ

Table 5

Tuble 0	
Errors between GPS survey and topog	graphic map data.

Point	Latitude (°)	Longitude (°)	Elevation (m) (GPS)	Elevation (m) (topographic map)	Error (m)
1	37.5462	101.7810	3972.0	3977.1	5.1
2	37.5371	101.8011	4137.5	4140.0	2.5
3	37.5403	101.8041	4009.6	4010.0	0.4
4	37.5398	101.8145	3882.1	3880.0	-2.1
5	37.5397	101.8146	3881.7	3883.5	1.8
6	37.5226	101.8369	4165.0	4163.0	-2.0
7	37.5308	101.8381	3935.4	3940.0	4.6
8	37.5210	101.8382	4156.8	4168.8	12.0
9	37.5439	101.8521	3710.2	3720.0	9.8

can probably be attributed to the increase in air temperature. Thus, combined with the above analysis, we also believe that glacier shrinkage and thinning in this area is mainly caused by the increase in summer temperature and FLH during the period 1972–2010.

Rate of glacier variation is controlled not only by climate change but also by topographic setting (e.g. aspect and slope), especially for glaciers at the regional scale or in the same climate condition (Scherler et al., 2011; Venkatesh et al., 2013). The aspects of these nine glaciers are primarily north. Therefore, the difference in thinning rate cannot be well explained by the aspect individually. The mean slope in the GPSmeasured area of each glacier is also calculated (Table 4). Generally, a higher mean thinning rate corresponded to a lower mean slope. In addition, the changes in thinning rate and slope for each glacier along paths are shown in Fig. 3b. The figure also shows a slightly negative relationship, with slope decreasing while thinning increases at each glacier. However, the high uncertainty of the vertical makes it of doubtful reliability for evaluating and discussing the correlation.

Glacier retreat and thinning in other mountains

Many studies have shown that the TP glaciers have responded to rapid warming by decreasing in area and losing ice mass in recent decades, with varying shrinkage statuses from region to region. For example, glacier mean area decreased by 0.57% yr⁻¹ in the southeastern TP, 0.41% yr⁻¹ in the central Himalayas, 0.42% yr⁻¹ in the western Himalayas, 0.29% yr⁻¹ in the northeastern TP, and 0.07% yr⁻¹ in the northwestern TP (Yao et al., 2012). In the southeastern TP, the Hailuogou glacier of the Gongga Mountains experienced a high thinning rate (1.1 m yr⁻¹ between 1966 and 2009) (Zhang et al., 2010), and Parlung No. 12 glacier also shows a significantly negative mass balance (-1.7 m yr^{-1} from 2005 to 2009) (Yao et al., 2012). Shangguan et al. (2010) found a thinning rate of about 0.4 m yr⁻¹ for two glaciers along the Yanglonghe river between 1956 and 2007, which is similar



Figure 4. Changes in annual precipitation (*P*), summer (June–August) air temperature (*T*) at the Menyuan weather station (2924 m), freezing-level height (*FLH*) at the Xining radiosonde station (2295 m) and the glacier area in the study area.

to the 0.41 m yr⁻¹ found in Dongkemadi Ice Field from 1961 to 2000 (Li et al., 2011) and slightly lower than rates in our study area (0.64 m yr⁻¹). However, Glacier No. 01 in the Urumqi river source of the Tien Shan had a much lower thinning rate of 0.24 m yr⁻¹ from 1962 to 2003 (Ye et al., 2005).

Conclusion

GPS is an efficient and accurate method for extracting data for glacier mass balance change. The nine glaciers in the NC and SG river source lost 9.9% of their total area in the period 1972–2010. The mean terminal altitude rose 39.3 m and glacier length retreated at a rate of 4.7 m yr⁻¹. The mean surface elevation decreased by 0.64 m yr⁻¹. Total glacier mass loss was at least 90.8 \pm 39.7 \times 10⁻³ km³ w.e. (0.1 \pm 0.05 m³/s) for 1972–2010 in NC and SG river source glaciers. The glaciers in the study area and in other mountains are retreating, decreasing in area and thinning in response to climate change during the past several decades.

Acknowledgments

We thank G. Zhang, B. Huang, F. Liu, D. Chen and F. Li for their hard field work. This paper has benefited from valuable comments and constructive suggestions by F. Chen and two anonymous reviewers, whose efforts are gratefully acknowledged. This study is financially supported by the National Basic Work Program of the Ministry of Science and Technology of China (Grant No. 2013FY111400), the National Natural Science Foundation of China (Grant Nos. 91125008 and 41171063), and the Program for New Century Excellent Talents in University (Grant No. NCET-13-0263).

References

- Aizen, V.B., Kuzmichenok, V.A., Surazakov, A.B., Aizen, E.M., 2007. Glacier changes in the Tien Shan as determined from topographic and remotely sensed data. Global and Planetary Change 56, 328–340.
- Anderson, B., Lawson, W., Owens, I., 2008. Response of Franz Josef Glacier Ka Roimata o Hine Hukatere to climate change. Global and Planetary Change 63, 23–30.
- Barrand, N.E., James, T.D., Murray, T., 2010. Spatio-temporal variability in elevation changes of two high-Arctic valley glaciers. Journal of Glaciology 56, 771–780.
- Berthier, E., Toutin, T., 2008. SPOT5-HRS digital elevation models and the monitoring of glacier elevation changes in North-West Canada and South-East Alaska. Remote Sensing of Environment 112, 2443–2454.
- Berthier, E., Arnaud, Y., Baratoux, D., Vincent, C., Rémy, F., 2004. Recent rapid thinning of the "Mer de Glace" glacier derived from satellite optical images. Geophysical Research Letters 31, L17401.
- Bolch, T., 2007. Climate change and glacier retreat in northern Tien Shan (Kazakhstan/ Kyrgyzstan) using remote sensing data. Global and Planetary Change 56, 1–12.
- Bolch, T., Pieczonka, T., Benn, D.I., 2011. Multi-decadal mass loss of glaciers in the Everest area (Nepal Himalaya) derived from stereo imagery. The Cryosphere 5, 349–358.
- Chen, F., Yu, Z., Yang, M., Ito, E., Wang, S., Madsen, D.B., Huang, X., Zhao, Y., Sato, T., John, B., Birks, H., Boomer, I., Chen, J., An, C., Wünnemann, B., 2008. Holocene moisture evolution in arid central Asia and its out-of-phase relationship with Asian monsoon history. Quaternary Science Reviews 27, 351–364.
- Ding, Y., Liu, S., Li, J., Shangguan, D., 2006. The retreat of glaciers in response to recent climate warming in western China. Annals of Glaciology 43, 97–105.
- Elizabeth, J.P., Manuel, H.-P., Paolo, S., Philip, M., Matt, A.K., 2011. A review of higher order ionospheric refraction effects on dual frequency GPS. Surveys in Geophysics 32, 197–253.

- Gao, H., He, X., Ye, B., Pu, J., 2012. Modeling the runoff and glacier mass balance in a small watershed on the Central Tibetan Plateau, China, from 1955 to 2008. Hydrological Processes 26, 1593–1603.
- Gardner, A.S., Moholdt, G., Cogley, J.G., Wouters, B., Arendt, A.A., Wahr, J., Berthier, E., Hock, R., Pfeffer, W.T., Kaser, G., Ligtenberg, S.R.M., Bolch, T., Sharp, M.J., Hagen, J.O., van den Broeke, M.R., Paul, F., 2013. A reconciled estimate of glacier contributions to sea level rise: 2003 to 2009. Science 340, 852–857.
- Hall, D.K., Bayr, K.J., Schöner, W., Bindschadler, R.A., Chien, J.Y.L., 2003. Consideration of the errors inherent in mapping historical glacier positions in Austria from the ground and space (1893–2001). Remote Sensing of Environment 86, 566–577.
- Howat, I.M., Joughin, I., Scambos, T.A., 2007. Rapid changes in ice discharge from Greenland outlet glaciers. Science 315, 1559–1561.
- Huss, M., 2013. Density assumptions for converting geodetic glacier volume change to mass change. The Cryosphere 7, 877–887.
- Jacob, T., Wahr, J., Pfeffer, W.T., Swenson, S., 2012. Recent contributions of glaciers and ice caps to sea level rise. Nature 482, 514–518.
- Kargel, J.S., Abrams, M.J., Bishop, M.P., Bush, A., Hamilton, G., Jiskoot, H., Kääb, A., Kieffer, H.H., Lee, E.M., Paul, F., Rau, F., Raup, B., Shroder, J.F., Soltesz, D., Stainforth, D., Stearns, L., Wessels, R., 2005. Multispectral imaging contributions to global land ice measurements from space. Remote Sensing of Environment 99, 187–219.
- Kaser, G., Cogley, J.G., Dyurgerov, M.B., Meier, M.F., Ohmura, A., 2006. Mass balance of glaciers and ice caps: consensus estimates for 1961–2004. Geophysical Research Letters 33, L19501.
- Kaser, G., Großhauser, M., Marzeion, B., 2010. Contribution potential of glaciers to water availability in different climate regimes. Proceedings of the National Academy of Sciences 107, 20223–20227.
- Keutterling, A., Thomas, A., 2006. Monitoring glacier elevation and volume changes with digital photogrammetry and GIS at Gepatschferner glacier, Austria. International Journal of Remote Sensing 27, 4371–4380.
- King, M., 2004. Rigorous GPS data-processing strategies for glaciological applications. Journal of Glaciology 50, 601–607.
- Kunz, M., King, M.A., Mills, J.P., Miller, P.E., Fox, A.J., Vaughan, D.G., Marsh, S.H., 2012. Multi-decadal glacier surface lowering in the Antarctic Peninsula. Geophysical Research Letters 39, L19502.
- Lemke, P., Ren, J., Alley, R.B., Allison, I., Carrasco, J., Flato, G., Fujii, Y., Kaser, G., Mote, P., Thomas, R.H., Zhang, T., 2007. Observations: changes in snow, ice and frozen ground. In: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K.B., Tignor, M., Miller, H.L. (Eds.), Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Li, J., Liu, S., Shangguan, D., Zhang, Y., 2010. Identification of ice elevation change of the Shuiguan River No. 4 glacier in the Qilian Mountains, China. Journal of Mountain Science 7, 375–379.
- Li, Z., Xing, Q., Liu, S., Zhou, J., Huang, L., 2011. Monitoring thickness and volume changes of the Dongkemadi Ice Field on the Qinghai-Tibetan Plateau (1969–2000) using Shuttle Radar Topography Mission and map data. International Journal of Digital Earth 1–17.
- Liu, C., Song, G., jin, M., 1992. Recent change and trend prediction of glaciers in Qilian Mountain. Memoirs of Lanzhou Institute of Glaciology and Geocryology Chinese Academy of Sciences, No. 7 1–9 (In Chinese).
- Liu, S., Sun, W., Shen, Y., Li, G., 2003. Glacier changes since the Little Ice Age maximum in the western Qilian Shan, northwest China, and consequences of glacier runoff for water supply. Journal of Glaciology 49, 117–124.
- Liu, S., Ding, Y., Shangguan, D., Zhang, Y., Li, J., Han, H., Wang, J., Xie, C., 2006. Glacier retreat as a result of climate warming and increased precipitation in the Tarim river basin, northwest China. Annals of Glaciology 43, 91–96.
- Meier, M.F., Dyurgerov, M.B., Rick, U.K., O'Neel, S., Pfeffer, W.T., Anderson, R.S., Anderson, S.P., Glazovsky, A.F., 2007. Glaciers dominate eustatic sea-level rise in the 21st century. Science 317, 1064–1067.
- Oerlemans, J., 2005. Extracting a climate signal from 169 glacier records. Science 308, 675–677.
- Ohmura, A., 2011. Observed mass balance of mountain glaciers and Greenland ice sheet in the 20th century and the present trends. Surveys in Geophysics 32, 537–554.
- Pan, B., Cao, B., Wang, J., Zhang, G., Zhang, C., Hu, Z., Huang, B., 2012. Glacier variations in response to climate change from 1972 to 2007 in the western Lenglongling mountains, northeastern Tibetan Plateau. Journal of Glaciology 58, 879–888.
- Paul, F., Kääb, A., Maisch, M., Kellenberger, T., Haeberli, W., 2004. Rapid disintegration of alpine glaciers observed with satellite data. Geophysical Research Letters 31, L21402.
- Paul, F., Barrand, N.E., Baumann, S., Berthier, E., Bolch, T., Casey, K., Frey, H., Joshi, S.P., Konovalov, V., Le Bris, R., Molg, N., Nosenko, G., Nuth, C., Pope, A., Racoviteanu, A., Rastner, P., Raup, B., Scharrer, K., Steffen, S., Winsvold, S., 2013. On the accuracy of glacier outlines derived from remote-sensing data. Annals of Glaciology 54, 171–182.
- Racoviteanu, A., Williams, M., Barry, R., 2008. Optical remote sensing of glacier characteristics: a review with focus on the Himalaya. Sensors 8, 3355–3383.

- Racoviteanu, A., Paul, F., Raup, B., Khalsa, S.J.S., Armstrong, R., 2009. Challenges and recommendations in mapping of glacier parameters from space: results of the 2008 Global Land Ice Measurements from Space (GLIMS) workshop, Boulder, Colorado, USA. Annals of Glaciology 50, 53–69.
- Radić, V., Hock, R., 2011. Regionally differentiated contribution of mountain glaciers and ice caps to future sea-level rise. Nature Geoscience 4, 91–94.
- Raup, B.H., Kääb, A., Kargel, J.S., Bishop, M.P., Hamilton, G., Lee, E., Paul, F., Rau, F., Soltesz, D., Khalsa, S.J.S., Beedle, M., Helm, C., 2007. Remote sensing and GIS technology in the Global Land Ice Measurements from Space Project. Computers & Geosciences 33, 104–125.
- Rignot, E., Rivera, A., Casassa, G., 2003. Contribution of the Patagonia icefields of South America to sea level rise. Science 302, 434–437.
- Rivera, A., Casassa, G., 1999. Volume changes on Pio XI glacier, Patagonia: 1975–1995. Global and Planetary Change 22, 233–244.
- Rivera, A., Casassa, G., Bamber, J., Kääb, A., 2005. Ice-elevation changes of Glaciar Chico, southern Patagonia, using ASTER DEMs, aerial photographs and GPS data. Journal of Glaciology 51, 105–112.
- Scherler, D., Bookhagen, B., Strecker, M.R., 2011. Hillslope–glacier coupling: the interplay of topography and glacial dynamics in high Asia. Journal of Geophysical Research: Earth Surface 116, F02019. http://dx.doi.org/10.1029/2010JF001751.
- Shangguan, D., Liu, S., Ding, Y., Ding, L., Xiong, L., Cai, D., Li, G., Lu, A., Zhang, S., Zhang, Y., 2006. Monitoring the glacier changes in the Muztag Ata and Konggur mountains, east Pamirs, based on Chinese glacier inventory and recent satellite imagery. Annals of Glaciology 43, 79–85.
- Shangguan, D., Liu, S., Ding, Y., Zhang, Y., Li, J., Li, X., Wu, Z., 2010. Changes in the elevation and extent of two glaciers along the Yanglonghe river, Qilian Shan, China. Journal of Glaciology 56, 309–317.
- Shi, Y., Liu, S., 2000. Estimation on the response of glaciers in China to the global warming in the 21st century. Chinese Science Bulletin 45, 668–672.
- Silverio, W., Jaquet, J.-M., 2005. Glacial cover mapping (1987–1996) of the Cordillera Blanca (Peru) using satellite imagery. Remote Sensing of Environment 95, 342–350.
- Surazakov, A.B., Aizen, V.B., 2006. Estimating volume change of mountain glaciers using SRTM and map-based topographic data. IEEE Transactions on Geoscience and Remote Sensing 44, 2991–2995.
- Venkatesh, T.N., Kulkarni, A.V., Srinivasan, J., 2013. Relative effect of slope and equilibrium line altitude on the retreat of Himalayan glaciers. The Cryosphere 6, 301–311.
- Wang, Z., Liu, C., You, G., Pu, J., Yang, H., Tian, P., 1981. Glacier Inventory of China I. Qilian Mountains. Science Press. Academia Sinica, Lanzhou Institute of Glaciology and Cryopedology, Beijing (In Chinese).
- Wang, J., Wang, J., Lu, C., 2003. Problem of coordinate transformation between WGS-84 and BEIJING 54. Journal of Geodesy and Geodynamics 23, 70–73 (in Chinese).
- Wang, N., He, J., Xi, J., Song, G., Pu, J., Wu, X., Chen, L., 2009. Study on the zone of maximum precipitation in the North slopes of central Qilian Mountains. Journal of Glaciology and Geocryology 31, 395–403 (In Chinese).
- Williams, R.S., Hall, D.K., Sigurdsson, O., Chien, J.Y.L., 1997. Comparison of satellite-derived with ground-based measurements of the fluctuations of the margins of Vatnajökull, Iceland, 1973–92. Annals of Glaciology 24, 72–80.
- Xie, Z., Wu, G., Wang, L., 1985. Recent advance and retreat of glaciers in Qilian Mountains. Memoirs of Lanzhou Institute of Glaciology and Geocryology Chinese Academy of Sciences, No. 5 82–90 (In Chinese).
- Xie, Z., ShangGuan, D., Zhang, S., Ding, Y., Liu, S., 2013. Index for hazard of glacier lake outburst flood of Lake Merzbacher by satellite-based monitoring of lake area and ice cover. Global and Planetary Change 107, 229–237.
- Yao, T., Wang, Y., Liu, S., Pu, J., Shen, Y., Lu, A., 2004. Recent glacial retreat in high Asia in China and its impact on water resource in Northwest China. Science in China Series D: Earth Sciences 47, 1065–1075.
- Yao, T., Thompson, L., Yang, W., Yu, W., Gao, Y., Guo, X., Yang, X., Duan, K., Zhao, H., Xu, B., Pu, J., Lu, A., Xiang, Y., Kattel, D.B., Joswiak, D., 2012. Different glacier status with atmospheric circulations in Tibetan Plateau and surroundings. Nature Climate Change 2, 663–667.
- Ye, B., Yang, D., Jiao, K., Han, T., Jin, Z., Yang, H., Li, Z., 2005. The Urumqi River source Glacier No. 1, Tianshan, China: changes over the past 45 years. Geophysical Research Letters 32, L21504.
- Ye, Q., Kang, S., Chen, F., Wang, J., 2006. Monitoring glacier variations on Geladandong mountain, central Tibetan Plateau, from 1969 to 2002 using remote-sensing and GIS technologies. Journal of Glaciology 52, 537–545.
- Zhang, Y., Fujita, K., Liu, S., Liu, Q., Wang, X., 2010. Multi-decadal ice-velocity and elevation changes of a monsoonal maritime glacier: Hailuogou glacier, China. Journal of Glaciology 56, 65–74.
- Zhang, S., Ye, B., Liu, S., Zhang, X., Hagemann, S., 2012. A modified monthly degree-day model for evaluating glacier runoff changes in China. Part I: model development. Hydrological Processes 26, 1686–1696.